

REFERENCES

- Aboul-Eid, H.Z. 1972. Pathogenicity of *Longidorus siddiqii* on Egyptian cotton (*Gossypium barbadense*). Plant Disease Reporter 56: 699-700.
- Aon, M.A. & Colaneri, A.C. 2001. Temporal and spatial evolution of enzymatic activities and physico-chemical properties in an agricultural soil. Applied Soil Ecology 18: 255-270.
- Atkinson, G.F. 1892. Some diseases of cotton. Alabama Agricultural Experiment Station, Auburn University Bulletin 41.
- Baird, R.E., Nankam, C., Moghaddam, P.F. & Pataky, J.K. 1994. Evaluation of seed treatments on shrunken-2 sweet corn. Plant Disease 78: 817-821.
- Barker, K.R. & Koennig, S.R. 1998. Developing sustainable systems for nematode management. Annual Review of Phytopathology 36: 165-205.
- Barker, K.R., Schmitt, D.P. & Campos, V.P. 1982. Response of peanut, corn, tobacco, and soybean to *Criconebella ornata*. Journal of Nematology 14: 576-581.
- Been, T.H. & Schomaker, C.H. 2006. Distribution patterns and sampling. Pages 302-324 in: Plant Nematology. R.N. Perry & M. Moens, eds. CABI, Wallingford, United Kingdom.
- Bergeson, G.B. & Ferris, J.M. 1986. Influence of tillage methods on *Pratylenchus spp.* for two soil types. Plant Disease 70: 326-328.
- Bolton, H., Elliot, L.F., Papendick, R.I. & Bezdicek, D.F. 1985. Soil microbial biomass and selected soil enzyme activities: Effect of fertilization and cropping practices. Soil Biology and Biochemistry 17: 297-302.

- Bridge, J. 1987. Control strategies in subsistence agriculture. Pages 389-420 in: Principles and practice of nematode control in crops. R.H. Brown & B.R. Kerry, eds. Academic Press, Australia.
- Buyer, J.S., Drinkwater, L.E. 1997. Comparison of substrate utilisation assay and fatty acid analyses of soil microbial communities. *Journal of Microbiological Methods* 30: 3-11.
- Chambers, K.R. 1987a. Epidemiology of maize root rot in South Africa. *Journal of Phytopathology* 118: 84-93.
- Chambers K.R. 1987b. Ability of fungal isolates from maize and sorghum to infect roots and reduce seedling emergence of two maize hybrids. *Plant Disease* 71:736-739.
- Channon, P. & Farina M.P.W. 1991. Are soil-borne diseases depressing yields of continuously-grown maize in Natal? *South African Journal of Plant and Soil* 8: 141-145.
- Conn, K.L., Tenuta, M. & Lazarovits, G. 2005. Liquid swine manure can kill *Verticillium dahliae* microsclerotia in soil by volatile fatty acid, nitrous acid, and ammonia toxicity. *Phytopathology* 95: 28-35.
- Deacon, J.W. & Scott, D.B. 1983. *Phialophora zeicola* sp. nov., and its role in the root rot-stalk rot complex of maize. *Transactions of the British Mycological Society* 81: 247-262.
- Deep I.W. & Lipps, P.E. 1996. Recovery of *Pythium arrhenomanes* and its virulence to corn. *Crop Protection* 15: 85-90.
- Demeure, Y., Freckman, D.W. & Van-Gundy, S D. 1979. In vitro response of four species of nematodes to desiccation and discussion of this and related phenomena. *Revue de Nématologie* 2: 203-210.

- Deng, S.P. & Tabatabai, M.A. 1997. Effects of crop residue management on soil enzyme activities in soils. III. Phosphatases and arylsulphatase. *Biology and Fertility of Soils* 24: 141-146.
- Denman, S., Knox-Davies, P.S., Calitz, F.J. & Lamprecht, S.C. 1995. Pathogenicity of *Pythium irregulare*, *P. sylvaticum* and *P. ultimum* to lucerne (*Medicago sativa*). *Australasian Plant Pathology* 24: 137-143.
- De Fede, K.L., Panaccione, D.G. and Sexstone, A.J. 2001. Characterization of dilution enrichment cultures obtained from size-fractionated soil bacteria by BIOLOG[®] community-level physiological profiles and restriction analysis of 16S rRNA genes. *Soil Biology and Biochemistry* 33: 1555-1562.
- De Waele, D. & Jordaan, E.M. 1988. Plant-parasitic nematodes on field crops in South Africa. 1. Maize. *Revue de Nématologie* 11: 65-74.
- Dick, R.P. 1984. Influence of long-term tillage and crop combinations on soil enzyme activities. *Soil Science Society of American Journal* 48: 569-574.
- Dick, R.P. 1992. A review: Long-term effects of agricultural systems on biochemical and microbial parameters. *Agriculture, Ecosystems and Environment* 40: 25-36.
- Dick, R.P. 1994. Soil enzyme activities as indicators of soil quality. Pages 107-124 in: *Defining soil quality for a sustainable environment*. J.W. Doran, D.C. Coleman, D.F. Bezdicek, & B.A. Stewart eds. SSSA Special Publication 35, Soil Science Society of America, Madison, WI.
- Dick, R.P. 1997. Soil enzyme activities as integrative indicators. Pages 121-156 in: *Biological indicators of soil health*. C.E. Pankhurst, B.M. Doube, & V.V.S.R. Gupta eds. CAB International, New York.

- Dick, R.P., Breakwell, D.P., Turco, R.F. 1996. Soil enzyme activities and biodiversity measurements as integrative microbiological indicators. In: Doran, J.W., Jones, A.J. (Eds.), Methods for assessing soil quality, SSSA Special Publication 49, Soil Science Society of America, Madison, WI, pp 247-271.
- Dorrance, A.E., Berry, S.A., Boven, P. & Lipps, P.E. 2004. Characterization of *Pythium* spp. from three Ohio fields for pathogenicity on corn and soybean and metalaxyl sensitivity. Plant Health Progress 2004: 1-7.
- Du Plessis, J. 2003. Maize production. Department of Agriculture, Pretoria, South Africa.
- Du Toit, J.J. 1968. Root rot of young maize plants – The causal fungi. South African Journal of Agricultural Science 11: 595-604.
- Du Toit, L.J., Kirby, H.W. & Pedersen, W.L. 1997. Evaluation of an aeroponics system to screen maize genotypes for resistance to *Fusarium graminearum* seedling blight. Plant Disease 81: 175-179.
- Eisenback, J.D. 1998. Glossary of plant nematology and related terms. CABI Publishing, Wallingford, United Kingdom.
- Elad, Y., Zvieli, Y. & Chet, I. 1986. Biological control of *Macrophomina phaseolina* (Tassi) Goid. by *Trichoderma harzianum*. Crop Protection 5: 288-292.
- Eno, C.F., Blue, W.G. & Good, J.M. 1955. The effect of anhydrous ammonia on nematodes, fungi, bacteria, and nitrification in some Florida soils. Soil Science Society of America Proceedings 19: 55-58.
- Farina, M.P.W. 2000. Don't ignore rotations. Subchapter 3.3.3, pages 1-7 in: A guide to no-till crop production in KwaZulu-Natal. A.Venter ed. ISBN 0-621-30095-0.

- Farina, M.P.W. & Channon, P. 1988. Acid-subsoil amelioration. 1. A comparison of several mechanical procedures. *Soil Science Society of America Journal* 52: 169-174.
- Flett, B.C., McLaren, N.W. & Wehner, F.C. 1998. Incidence of ear rot pathogens under alternating corn tillage practices. *Plant Disease* 82: 781-784.
- Furstenberg, J.P. & Heyns, J. 1978. The effect of cultivation on nematodes. Part 1. *Rotylenchulus parvus*. *Phytophylactica* 10: 77-80.
- Garbeva, P., Van Veen, J.A., Van Elsas, J.D. 2004. Microbial diversity in soil: Selection of microbial populations by plant and soil type and implications for disease suppressiveness. *Annual Review of Phytopathology*. 42: 243-270.
- Garcia, C., Hernandez, T & Costa, F. 1994. Microbial activity in soils under Mediterranean environmental conditions. *Soil Biology and Biochemistry* 26: 1185-1191.
- Garcia, C., Hernandez, T., Roldan, A & Martin, A. 2002. Effect of plant cover decline on chemical and microbiological parameters under Mediterranean climate. *Soil Biology and Biochemistry* 34: 635-642.
- Garland, J.L. 1996. Patterns of potential C source utilization by rhizosphere communities. *Soil Biology and Biochemistry* 28: 223-230.
- Garland, J.L. & Mills, A.L. 1991. Classification and characterization of heterotrophic microbial communities on the basis of patterns of community-level sole-carbon-source utilization. *Applied Environmental Microbiology* 57: 2351-2359.
- Glazer, I. & Orion, D. 1983. Studies on anhydrobiosis of *Pratylenchus thornei*. *Journal of Nematology* 1983; 15: 333-338.

- Griffiths, B.S., Ritz, K. & Wheatley, R.E. 1994. Nematodes as indicators of enhanced microbiological activity in a Scottish organic farming system. *Soil Use and Management* 10: 20-24.
- Graham, R.D. & Webb, M.J. 1991. Micronutrients and disease resistance and tolerance in plants. Pages 329-371 in: *In Micronutrients in Agriculture*, 2nd ed, SSSA Book Series, no.4, Madison WI, USA.
- Havlin, J.I., Beaton, J.D., Tisdale, S.L & Nelson, W.R. 1999. *Soil Fertility and Fertilizers*. Pages 86-153. Prentice Hall, Incorporated, Upper Saddle River, NJ.
- Haycock, P.P.J., Woods, S.R., Grove, I.G. & Hare, M.C. 2006. Chemical control of nematodes. Pages 392-410 in: *Plant Nematology*. R.N. Perry & M. Moens, eds. CABI, Wallingford, United Kingdom.
- Hellinga, J.H., Bouwman, J.J., Scholte, K. & s'Jacob, J.J. 1983. Causes of root rot in maize on sandy soil. *Netherlands Journal of Plant Pathology* 89: 229-237.
- Hornby, D. & Ullstrup, A.J. 1967. Fungal populations associated with maize roots. Composition and comparisons of mycofloras from genotypes differing in root rot resistance. *Phytopathology* 57: 869-875.
- Howard, D.D., Chambers, A.Y. & Lessman, G.M. 1998. Rotation and fertilization effects on corn and soybean yields and soybean cyst nematode populations in a no-tillage system. *Agronomy Journal* 90: 518-522.
- Huber, D.M. 1989. The role of nutrition in the take-all disease of wheat and other small grains. Pages 46-74 in: *Soilborne Plant Pathogens. Management of Diseases with Macro-and Micro-elements*. A.W. Engelhard ed. APS Press, St Paul, Minnesota.

- Huber, D.M. 1991. The use of fertilizers and organic amendments in the control of plant diseases. Pages 405-494 in: Handbook of Pest Management in Agriculture, vol. 1, 2nd ed, D, Pimentel (ed), CRC Press, Boca Raton, FL.
- Inomoto, M.M., Motta, L.C.C., Machado, A.C.Z. & Sazaki, C.S.S. 2006. Reaçãp de dez cobeturas vegetais a *Pratylenchus brachyurus*. Nematologia Brasileira 30: 151-157.
- Jackson, T. 2005. Managing corn nematodes common to Nebraska. Crop Watch July 2005. [<http://cropwatch.unl.edu>].
- Johnson, A.W. 1975. Resistance of sweet corn cultivars to plant-parasitic nematodes. Plant Disease Reporter 59: 373-376.
- Jordan, D., Kremner, R.J., Bergfield, W.A., Kim, K.Y. & Cacnio, V.N. 1995. Evaluation of microbial methods as potential indicators of soil quality in historical agricultural fields. Biology and Fertility of Soils 19: 297-302.
- Kandeler, E., Gerber, H. 1988 Short-term assay of soil urease activity using colorimetric determination of ammonium. Biology and Fertility of Soils 6: 68 – 72.
- Kandji, S.T., Ogol, C.K.P.O. & Albrecht, A. 2003. Crop damage by nematodes in improved-fallow fields in western Kenya. Agroforestry Systems 57: 51-57.
- Khan, S.W. 1970. Enzymatic activity in a gray wooded soil as influenced by cropping systems and fertilizers. Soil Biology and Fertility 2: 137-139.
- Keetch, D.P. 1989. A perspective of plant nematology in South Africa. South African Journal of Science 85: 506-508.

- Keetch, D.P. & Milne, D.L. 1982. The control of plant-parasitic nematodes. Pages 113-129 in: Nematology in Southern Africa. D.P. Keetch & J. Heyns, eds. Science Bulletin No. 400. Department of Agriculture and Fisheries, Pretoria, South Africa.
- Kleynhans, K.P.N. 1997. Collecting and preserving nematodes. Manual for a SAFRINET course in practical Nematology. SAFRINET, the Southern African (SADC) LOOP of BioNet-INTERNATIONAL, Pretoria.
- Kleynhans, K.P.N., Van Den Berg, E., Swart, A., Marais, M. & Buckley, N.H. 1996. Plant nematodes in South Africa. Plant Protection Research Institute Handbook No. 8. ARC-Plant Protection Research Institute, Pretoria.
- Krüger, W. 1970. Würzel – und stammfaule bei mais. II. Wurzelfaule verunsackerend organismen im ‘maiz-dreieck’ Sudafrikas. *Phytopathologische Zeitschrift* 67: 345-351.
- Krüger, W. & Speakman, J.B. 1997. Crop rotation studies with regard to root and stem base rots in wheat and root rots in maize. *Mitteilungen aus der Biologischen Bundesanstalt für Land und Forstwirtschaft Berlin Dahlem* 261:129 (Abstr.).
- Kumar, M & Agarwal, V.K. 1998. Effect of fungicidal seed treatment on seedling vigour of maize. *Seed Research* 26: 147-151.
- Lamprecht, S. C. 2007. Effect of crop rotation on crown and root rot severity of maize in rotation systems at Vaalharts. Report to ARC-GCI.
- Lamprecht, S.C., Farina, M.P.W., Habig, J.H., Thibaud, G.R., & Bloem, J.F. 2006. The role and importance of soilborne diseases in yield depression of maize. Report to No-till club, KZN, and Maize Trust.
- Lipps, P.E. & Deep, I.W. 1991. Influence of tillage and crop rotation on yield, stalk rot and recovery of *Fusarium* and *Trichoderma* spp. from corn. *Plant Disease* 75: 828-833.

- Magurran, A.E. 1988. Ecological diversity and its measurement. Princeton University Press, Princeton, New Jersey, pp. 179.
- Marais, M. & Swart, A. 2007. Plant nematodes in South Africa. 8. Bizana, Lusikisiki and Port St Johns area, Eastern Cape Province. African Plant Protection 13: 16-27.
- Mannering, J.V. & Griffiths, D.P. 2000. Value of crop rotation under various tillage systems. Subchapter 3.3.1, pages 1-5 in: A guide to no-till crop production in KwaZulu-Natal. A. Venter ed. ISBN 0-621-30095-0.
- Marasas, W.F.O., Nelson, P.E. & Toussoun, T.A. 1984. Toxigenic *Fusarium* species: Identity and Mycotoxicology. The Pennsylvania State University Press, Pennsylvania.
- Martens, D.A., Johanson, J.B. & Frankenberger, W.T. Jr. 1992. Production and persistence of soil enzymes with repeated additions of organic residues. Soil Science 153: 53-61.
- Martin, G.C. Way, J.J. & Armstrong, A.M. 1975. Soil fumigation and the control of *Pratylenchus* spp. and other plant parasitic nematodes affecting maize production in Rhodesia. Paper presented at the second Symposium of the Nematological Society of Southern Africa, Stellenbosch, April 1975.
- McCormick, S. 2003. The role of DON in pathogenicity. Pages 165-183 in: Fusarium Head Blight of Wheat and Barley. K.L. Leonard and W.R. Bushell eds. APS Press, Minnesota.
- McDonald, A.H. & De Waele, D. 1987. Control strategies against nematode pests on maize. Proceedings of the 7th South African maize Breeding Symposium 1986. Technical Communication No 212. Department of Agriculture and Water Supply, Pretoria, South Africa.

- McDonald, A.H., Lui's, J.H. Loots, G.L. & De Waele, D. 1987. Chemical control of lesion nematodes (*Pratylenchus* spp.) on maize in South Africa. *Phytophylactica* 19: 479-483.
- McDonald, A.H. & Nicol, J.M. 2005. Nematode parasites of cereals. Pages 131-191 in: *Plant parasitic nematodes in subtropical and tropical agriculture*, second edition. M. Luc, R.A. Sikora & J. Bridge, eds. CABI Publishing, Wallingford, United Kingdom.
- McFadden, A.G. & Sutton, J.C. 1975. Relationships of populations of *Trichoderma* spp. in soil to disease in maize. *Canadian Journal of Plant Science* 55: 579-586.
- McLaughlin, A. & Mineau, P. 1995. The impact of agricultural practices on biodiversity. *Agriculture, Ecosystems & Environment* 55: 201-212.
- McSorley, R. & Gallaher, R.N. 1993. Effect of crop rotation and tillage on nematode densities in tropical corn. *Journal of Nematology (Supplement)* 25: 814-819.
- Miller, R.E. 1964. A root rot complex of field maize: Symptomatology, sequence of fungi involved and the effect of corn root exudation on these fungi. *Dissertation Abstracts* 24: 3921.
- Miller, M. & Dick, R.P. 1995a. Dynamics of soil C and microbial biomass on whole soil aggregates in two cropping systems. *Applied Soil Ecology* 2: 253-261.
- Miller, M. & Dick, R.P. 1995b. Thermal stability and activities of soil C enzymes as influenced by crop rotations. *Soil Biology and Biochemistry* 27: 1161-1166.
- Moreno-González, J., Andrés Ares, J.L., Alonso Ferro, R. & Campo Ramírez, L. 2004. Genetic and statistical models for estimating genetic parameters of maize seedling resistance to *Fusarium graminearum* Schwabe root rot. *Euphytica* 137: 55-61.

- Munkvold, G.P. & O'Mara, J.K. 2002. Laboratory and growth chamber evaluation of fungicidal seed treatments for maize seedling blight caused by *Fusarium* species. *Plant Disease* 86: 143-150.
- Nel, A., Krause, M. & Khelawanlall, N. 2003. A guide to the control of plant diseases. Department of Agriculture, Republic of South Africa.
- Netscher, C. & Seinhorst, J.W. 1969. Propionic acid better than acetic acid for killing nematodes. *Nematologica* 15: 286.
- Niblack, T. 2007 iTime to sample for nematodes in at-risk corn fields. *The Bulletin* 12, Article 6, June 15, 2007.
[<http://www.ipm.uiuc.edu/bulletin/article.php?issueNumber=12&issueyear=2007&article=6>].
- Olowe, R. & Corbett, D.C.M. 1976. Aspects of the biology of *Pratylenchus brachyurus* and *P. zaeae*. *Nematologica* 22: 202-211.
- Osborne, L., Smolik, J. & Osborne, S. 2004. Nematode community dynamics in no-till crop rotations for the northern Great Plains [Abstract]. Joint meeting of the ASA, CSSA, SSSA with the Canadian Society of Soil Science, October 31- November 4, 2004.
[http://www.ars.usda.gov/research/publications/publications.htm?SEQ_NO_115=16688].
- Palmer, L.T. & Kommedahl, T. 1969. Root-infecting *Fusarium* species in relation to rootworm infestations in corn. *Phytopathology* 59: 1613-1617.
- Palmer, L.T. & McDonald, D.H. 1974. Interaction of *Fusarium* spp. and certain plant parasitic nematodes on maize. *Phytopathology* 64: 14-17.

- Palojärvi, A., Sharma, S., Rangger, A., Von Lutzow, M., Insam, H. 1997. Comparison of BIOLOG® and Phospholipid Fatty acid patterns to detect changes in microbial community. In: Insam, H., Rangger, A. (Eds.), *Microbial Communities – Functional versus Structural Approaches*. Springer-Verlag, Berlin, Heidelberg, Germany, pp. 37-48.
- Pankaj, S.H.K., Gaur, H.S. & Singh, A.K. 2006. Effect of zero tillage on the nematode fauna in a rice-wheat cropping system. *Nematologia Meditteranea* 34: 175-178.
- Pankhurst, C.E. & Lynch, J.M. 1995. The role of soil microbiology in sustainable intensive agriculture. *Advances in Plant Pathology*. 11: 229-247.
- Paterson, A. 2007. No-till farming. *The Witness*, Aug. 17.
- Pioli, R.N., Mozzoni, L. & Morandi, E.N. 2004. First report of pathogenic association between *Fusarium graminearum* and soybean. *Plant Disease* 88: 220.
- Ramsey, M.D. 1990. Etiology of root and stalk rots of maize in north Queensland. Disease development and associated fungi. *Australasian Plant Pathology* 19: 2-12.
- Rao, B. Schmitthenner, A.F., Caldwell, R. & Ellett, C.W. 1978. Prevalence and virulence of *Pythium* species associated with root rot of corn in poorly drained soil. *Phytopathology* 68: 1557-1563.
- Riekert, H.F. 1996a. Greenhouse assessment of maize growth yield response to nematode control with aldicarb. *African Crop Science Journal* 4: 471-475.
- Riekert, H.F. 1996b. Economic feasibility of nematode control in dryland maize in South Africa. *African Crop Science Journal* 4: 477-481.

- Ritz, K., McHugh, M. & Harris, J. 2003. Biological diversity and function in soils: contemporary perspectives and implications in relation to the formulation of effective indicators. Pages 1-11 in: OECD expert meeting on soil erosion and soil biodiversity indicators. Rome, March 2003.
- Roper, M.M. & Gupta, V. 1995. Management practices and soil biota. Australian Journal of Soil Research 33: 332-339.
- Schaafsma, A.W., Tamburic-Ilincic, L. & Hooker, D.C. 2005. Effect of previous crop, tillage, field size, adjacent crop, and sampling direction on airborne propagules of *Gibberella zeae/Fusarium graminearum* head blight severity and deoxynivalenol accumulation in winter wheat. Canadian Journal of Plant Pathology 27: 217-224.
- Scholte, K. 1987. Relationship between cropping frequency, root rot and yield of silage maize on sandy soils. Netherlands Journal of Agricultural Science 35: 473-486.
- Scott, D.B. 1982. Wortel- en stamvrot van mielies. Department of Agriculture and Fisheries, RSA, Technical Communication no. 118: 1-6.
- Scott, D.B., Kilian, W.H. & Miles, W.S. 1992. Influence of crop production practices on *Pythium* infections and yield of winter wheat in fumigated and non-fumigated soil. South African Journal of Plant and Soil 9: 14-18.
- Shepherd, J.A. 1977. Hosts of non-gall-forming nematodes associated with tobacco in Rhodesia. Rhodesian Journal of Agricultural Research 15: 95-97.
- Smit, E. 1998. Complex of root infecting fungi isolated from maize grown under various tillage practices. South African Journal of Plant and Soil 15: 116-120.

- Smit, E., Van Rensburg, G.D.J. & Rijkenberg, F.H.J. 1997. Number of isolates of maize root fungi in different crop rotation systems. *South African Journal of Plant and Soil* 14: 127-130.
- Srobarova, -A. & Eged, -S. 2005. *Trichoderma* and sulphoethyl glucan reduce maize root rot infestation and fusaric acid content. *Plant, Soil and Environment* 51: 322-327.
- Sumner, D.R. & Bell, D.K. 1982. Root diseases induced in corn by *Rhizoctonia solani* and *Rhizoctonia zea*. *Phytopathology* 72: 86-91.
- Sumner, D.R. & Dowler, C.C. 1983. Herbicide, planting date and root disease interactions in corn. *Plant Disease* 67: 513-517.
- Sumner, D.R., Hall, M.R., Gay, J.D., MacDonald, G. Savage, S.I. & Bramwell, R.K. 2002. Root diseases, weeds, and nematodes with poultry litter and conservation tillage in a sweet corn snap bean double crop. *Crop Protection* 21: 963-972.
- Taylor, D.P. 1961. Biology and host-parasite relationships of the spiral nematode, *Helicotylenchus microlobus*. *Proceedings of the Helminthological Society of Washington* 28: 60-66.
- Tenuta, M. & Lazarovits, G. 2002. Ammonia and nitrous acid from nitrogenous amendments kill the microsclerotia of *Verticillium dahliae*. *Phytopathology* 92: 255-264.
- Todd, T.C. 1991. Effects of cropping regime on populations of *Belonolaimus* sp. and *Pratylenchus scribneri* in sandy soil. *Journal of Nematology (Supplement)* 23: 646-651.
- Townshend, J. L. 1987. Anhydrobiosis in *Pratylenchus penetrans* and *Tylenchorhynchus* n.sp. in cultivated soils cropped to winter rye. *Journal of Nematology* 19: 164-171.

- Turner, B.L., Hopkins, D.W., Haygarth, P.M. & Ostle, N. 2002. Glucosidase activity in pasture soils. *Applied Soil Ecology* 20: 157-162.
- Unger, P.W. 1992. Conservation tillage systems. *Advanced Soil Science* 13: 27-68.
- Van den Berg, E. 1978. The genus *Rotylenchulus* Linford & Oliveira, 1940 (Rotylenchulinae: Nematoda) in South Africa. *Phytophylactica* 10: 57-64.
- Van Wyk, P.S., Scholtz, D.J. & Marasas, W.F.O. 1988. Protection of maize seedlings by *Fusarium moniliforme* against infection by *Fusarium graminearum* in soil. *Plant and Soil* 107: 251-257.
- Verstraete, W. and Voets, J.P. 1977. Soil microbial and biochemical characteristics in relation to soil management and fertility. *Soil Biology and Biochemistry* 9: 253-258.
- Walters, M.C. 1979. Present status of knowledge of nematode damage and control in South Africa. Technical Communication No 152. Department of Agricultural Technical Services, Pretoria, South Africa.
- Wang, H. & Davis, R.M. 1997. Susceptibility of selected corn cultivars to seedling disease pathogens and benefits of chemical seed treatments. *Plant Disease* 81: 1085-1088.
- Warren, H.L. & Kommedahl, T. 1973. Prevalence and pathogenicity to corn of *Fusarium* species from corn roots, rhizosphere, residues and soil. *Phytopathology* 63: 1288-1290.
- White, D.G. 1999. Compendium of corn diseases. 3rd Ed. APS Press, The American Phytopathology Society, Minnesota.

- Whitney, N.J. & Mortimore, C.G. 1961. Root and stalk rot of field corn in south-western Ontario. II. Development of the disease and isolation of organisms. *Canadian Journal of Plant Science* 41: 854-861.
- Williams, L.E. & Schmitthenner, A.F. 1963. Effect of crop rotation on yields, stalk and root rot of corn. *Phytopathology* 53: 1412-1414.
- Windham, G.L. 1998. Corn. Pages 335-357 in: Plant and nematode interactions. K.R.Barker, G.A. Pederson & G.L. Windham, eds. American Society of Agronomy, Crop Science Society of America, Soil Science Society of America, Madison, United States of America.
- Winding, A, Hendriksen, N.B. 1997. BIOLOG® substrate utilisation assay for metabolic fingerprints of soil bacteria: incubation effects. In: Insam, I., Rangger, A. (Eds.), *Microbial Communities: Functional versus Structural Approaches*. Springer-Verlag, Berlin, Heidelberg, Germany, pp. 195-205.
- Yassin, A.M. 1974. A note on *Longidorus* and *Xiphinema* species from the Sudan. *Nematologia Meditteranea* 2: 141-147.
- Yeates, G.W. & Bongers, T. 1995. Nematode diversity in agroecosystems. *Agriculture, Ecosystems and Environment* 74: 113-135.
- Young, T.R. & Kucharek, T.A. 1977. Successions of fungal communities in roots and stalks of hybrid field corn grown in Florida. *Plant Disease Reporter* 61: 76-80.
- Zhang, B.Q., Chen, W.D. & Yang, X.B. 1998. Occurrence of *Pythium* species in long-term maize and soybean monoculture and maize soybean rotation. *Mycological Research* 102: 1450-1452.